

Contents

696
ESSAY
Bureaucracy in general practice: time to halt a proliferating malaise
Edin Lakasing and Heather Francis

699
ESSAY
Family medicine in Japan
Patrick Hutt

702
ESSAY
The levels of medical effectiveness
Peter Davies

704
DIGEST
Book review
Peter D Toon

Book review
Roger Jones

705
DIGEST
Epidemics of epidemics
Mike Fitzpatrick

706
What next?
Helen Lester

NICE GUIDELINES ON COMPLEMENTARY/ALTERNATIVE MEDICINE: MORE CONSISTENCY AND RIGOUR ARE NEEDED

Recent NICE guidelines have recommended that patients suffering from persistent, non-specific low back pain should be offered 'one of the following treatment options, taking into account patient preference: an exercise programme, a course of manual therapy, or a course of acupuncture'.¹ No previous document by NICE has been so explicitly positive about complementary/alternative medicine (CAM). This, it seems, is a good occasion to review the totality of the recommendations from NICE guidelines in respect to CAM.

As of July 2009, NICE have published 88 clinical guidelines.² We have reviewed 83 of these (five have been replaced and were no longer available). The majority of these documents make no mention of CAM at all, often, it seems, for good reason; there are many conditions for which there is no evidence-based CAM option. Several guidelines refer to 'holistic' care, nutrition, vitamin therapy, and exercise, which some might view as CAM. For example, those relating to chronic and/or degenerative conditions explicitly recognise the need for holistic patient care and management, or focus on dietary and lifestyle interventions. A small number of the guidelines recognise various forms of CAM as an effective treatment. For instance, the depression guidelines state that St John's Wort is more effective than placebo. Yet the use of St John's wort is not recommended 'because of uncertainty about appropriate doses, variation in the nature of the preparations, and potential serious interactions with other drugs'.¹

Many of the guidelines which do incorporate CAM state or imply that patients themselves should determine the value of CAM by trial and error, or make statements such as 'some patients may find CAM useful'. The majority of these documents refer to CAM with the assertion that 'further research is needed' or that 'evidence is insufficient for firm recommendations'. In a number of instances (for example, guidelines relating to anxiety, type 2 diabetes, hypertension, and stroke) the impression is given that there is little evidence

to indicate that CAM approaches may be valuable, while reasonably sound data do exist.³ In other instances, the evidence for CAM is hopeful but the NICE verdict is discouraging. For example, the osteoarthritis guidelines state that 'results from acupuncture studies are mixed'. Yet several systematic reviews of acupuncture for osteoarthritis have drawn positive overall conclusions.⁴⁻⁶ Acupuncturists have therefore argued that this NICE guideline 'gives the reader a distinctly negative impression of acupuncture. This could be against patients' best interests ...'.⁷ The opposite scenario could be true for the recent back pain guidelines.¹ It has been argued that they over-estimated the effectiveness and under-estimated the risks of spinal manipulation.⁸

In conclusion, our analysis shows that many NICE guidelines have evaluated CAM. But there is a substantial degree of inconsistency: some are comprehensive, while others are not. The decision of whether or not to consider CAM for any given topic appears to be somewhat arbitrary. Generally speaking, this inconsistency seems to be at odds with NICE's excellent track record of evaluating conventional treatments, by utilising a transparent and rigid hierarchy of evidence for its recommendations. We therefore suggest that, in future, NICE should evaluate CAM by the same standards as conventional medicine.

Edzard Ernst and Rohini Terry

REFERENCES

1. Savigny P, Watson P, Underwood M. Early management of persistent non-specific low back pain: summary of NICE guidance. *BMJ* 2009; **338**: 1441-1442.
2. National Institute for Health and Clinical Excellence (NICE). Our guidance. <http://www.nice.org.uk/guidance/index.jsp> (accessed 14 Aug 2009).
3. Ernst E, Pittler MH, Wider B, Boddy K. *The desktop guide to complementary and alternative medicine*. 2nd edn. Edinburgh: Elsevier Mosby, 2006.
4. Kwon YD, Pittler MH, Ernst E. Acupuncture for peripheral joint osteoarthritis: a systematic review and meta-analysis. *Rheumatology (Oxford)* 2006; **45**: 1331-1337.
5. Ezzo J, Hadhazy V, Birch S, et al. Acupuncture for osteoarthritis of the knee: a systematic review. *Arthritis Rheum* 2001; **44**(4): 819-825.
6. White A, Foster NE, Cummings M, Barlas P. Acupuncture treatment for chronic knee pain: a systematic review. *Rheumatology (Oxford)* 2007; **46**(3): 384-390.
7. White A. NICE guideline on osteoarthritis: is it fair to acupuncture? No. *Acupunct Med* 2009; **27**(2): 70-72.
8. Ernst E. Spinal manipulation for the early management of persistent non-specific low back-pain — a critique of the recent NICE guidelines. *Int J Clin Pract* 2009: in press.

DOI: 10.3399/bjgp09X454232